



Universidad
Carlos III de Madrid



This is a postprint version of the following published document:

Barbero G., J. F., Prieto, J. & S.
Villaseñor, E. J. (2013): Band structure in
the polymer quantization of the harmonic
oscillator. *Classical and Quantum Gravity*,
30 (16), 165011.

DOI: [10.1016/j.conbuildmat.2015.04.060](https://doi.org/10.1016/j.conbuildmat.2015.04.060)

© IOP Publishing Ltd, 2013

Band structure in the polymer quantization of the harmonic oscillator

J Fernando Barbero G^{1,2}, Jorge Prieto³ and Eduardo J S Villaseñor^{2,3}

Abstract

We discuss the detailed structure of the spectrum of the Hamiltonian for the polymerized harmonic oscillator and compare it with the spectrum in the standard quantization. As we will see the non-separability of the Hilbert space implies that the *point* spectrum consists of bands similar to the ones appearing in the treatment of periodic potentials. This feature of the spectrum of the polymeric harmonic oscillator may be relevant for the discussion of the polymer quantization of the scalar field and may have interesting consequences for the statistical mechanics of these models.

1. Introduction

The polymer quantization of the harmonic oscillator has been considered as a toy model to gain a deeper understanding about some features of loop quantum gravity (LQG) [1–3]. It has also been used to discuss several relevant issues in loop quantum cosmology (LQC) [4, 5] and may be useful to study field theories such as the scalar field [6–10]. One of the consequences of the use of the more exotic, *non-separable*, Hilbert spaces characteristic of this approach is that the uniqueness theorem of Stone–von Neumann—in the case of quantum mechanical systems with a finite number of degrees of freedom—can be sidestepped and it is then possible to work with non-standard representations of the Weyl algebra (see [11] for the use of the polymer approach to discuss conceptual issues in quantum mechanics).

The available literature on this subject discusses some features of the spectrum of one of the infinitely many possible generalizations of the standard harmonic oscillator Hamiltonian that can be written in the new setting. The particular choice made in the literature gives a

¹ Instituto de Estructura de la Materia, CSIC, Serrano 123, E-28006 Madrid, Spain

² Grupo de Teorías de Campos y Física Estadística, Instituto Universitario Gregorio Millán Barbany, Universidad Carlos III de Madrid, Unidad Asociada al IEM-CSIC, Madrid, Spain

³ Instituto Gregorio Millán, Grupo de Modelización y Simulación Numérica, Universidad Carlos III de Madrid, Avda. de la Universidad 30, E-28911 Leganés, Spain

E-mail: fbarbero@iem.cfm.csic.es, jorgeprietoarranz@gmail.com and ejsanche@math.uc3m.es

well known differential equation—the Mathieu equation—for the eigenvalue problem and has led to the conclusion that the spectrum of the polymerized Hamiltonian is very similar to the standard one in quantum mechanics in the limit where the characteristic length parameter $q_0\ell$, that controls the quantum polymer harmonic oscillator Hamiltonian, becomes very small [2, 3, 12]. (Here, ℓ has dimension of length and q_0 is dimensionless.) There is, however, a disturbing issue that should be addressed. If the spectrum is a pure point one consisting of a countable set of non-degenerate eigenvalues (close to those of the standard quantized harmonic oscillator) then something must be missing because the countable set of eigenvectors cannot provide an orthonormal basis for the non-separable Hilbert space of the polymerized harmonic oscillator (any such orthonormal basis should be uncountable). As we show in the paper the actual spectrum of the Hamiltonian operator for the polymerized harmonic oscillator consists of an uncountable number of eigenvalues grouped in *bands*, very much like those appearing in the study of periodic potentials in standard quantum mechanics. The main difference between the spectrum of the polymerized harmonic oscillator and that of a standard periodic potential is that, in the former case, the points in the spectrum are actual eigenvalues (i.e. they belong to the point spectrum) associated with *normalizable* states whereas for the standard quantization of periodic potentials the bands correspond to the *continuous* spectrum. As a consequence of this, although some parts of the spectrum of the polymerized harmonic oscillator (in particular the lowest eigenvalues), resemble the lowest lying part of the standard harmonic oscillator spectrum—when the characteristic length $q_0\ell$ tends to zero—some crucial differences arise. These are associated, in particular, with the fact that in the ‘limit’ $q_0 \rightarrow 0$ the eigenvalues become infinitely degenerate. This has some far reaching consequences, in particular, for the statistical mechanics of these models because the standard definition for both the statistical entropy in the microcanonical ensemble and the partition function in the canonical one fail to be well defined. It may also be relevant for the polymer quantization of the scalar fields—that relies heavily upon the quantization of the harmonic oscillator.

The paper is organized as follows. After this introduction we briefly review in section 2 the main features of the Hilbert space of almost periodic functions on the real line $AP(\mathbb{R}, \mathbb{C})$ or, equivalently, $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$. We will emphasize, in particular, the fact that it is isomorphic to the Hilbert space $\ell^2(\mathbb{R})$ (the isomorphism realized in concrete terms by the Fourier–Bohr transform). We will quickly discuss in section 3 the position representation (PR) and momentum representation (MR) of the Weyl algebra in this framework. We will then present different ways of writing suitable Hamiltonians that mimic the standard Hamiltonian for the ordinary harmonic oscillator in the ‘limit’ in which certain parameter—that has to be introduced to approximate one of the quadratic pieces of the ordinary Hamiltonian—goes to zero. In section 4 we study in detail the spectrum of the resulting Hamiltonians in the non-separable space of polymerized quantum mechanics. By relying on mathematical theorems developed since the 1960s we will show that the spectra of these Hamiltonians can be obtained by considering the auxiliary problem of obtaining the spectrum of a Hamiltonian with a periodic potential in the separable Hilbert space $L^2(\mathbb{R}, \mu_{\mathbb{R}})$, where $\mu_{\mathbb{R}}$ is the Lebesgue measure. This is straightforward because the Bloch theorem (or the Floquet theory for differential equations with periodic coefficients) provides a complete solution. The main features of the band spectrum for periodic potentials have been extensively studied [13] and many rigorous, and even completely solvable, examples have been discussed in the literature. We briefly discuss some concrete examples of Hamiltonians for the polymerized harmonic oscillator in section 5. In addition to the popular Hamiltonian that leads to the Mathieu equation for the computation of the energy eigenvalues, we will also comment on another natural choice for the potential: a periodic extension of a purely quadratic potential defined on a symmetric interval about the origin with a periodicity controlled by the characteristic length parameter

$q_0\ell$. We will also comment briefly on the Lamé potentials in order to argue that the number of bands in the spectrum can actually be finite. The main reason to do this is to emphasize the enormous ambiguity present in the problem. We show in section 6 how the structure of the spectrum makes it difficult to define the partition function for these models and study their statistical mechanics [14], at least if one relies on standard energy-based ensembles and does not introduce extra conditions to eliminate unwanted parts of the spectrum. We end the paper in section 7 with our conclusions and comments regarding, in particular, the relevance of the banded structure of the spectrum for the physical applications of these models.

2. Polymer Hilbert spaces

The Hilbert spaces $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ and $AP(\mathbb{R}, \mathbb{C})$, that will be use in the paper, are unitarily isomorphic to the non-separable complex Hilbert space

$$\ell^2(\mathbb{R}) := \left\{ \Psi : \mathbb{R} \rightarrow \mathbb{C} : \sum_{x \in \mathbb{R}} |\Psi(x)|^2 < \infty \right\}$$

with scalar product

$$\langle \Psi_1, \Psi_2 \rangle_{\ell^2(\mathbb{R})} = \sum_{x \in \mathbb{R}} \overline{\Psi_1(x)} \Psi_2(x).$$

Notice that, as the sums extend over the whole real line, the set $\{x \in \mathbb{R} : \Psi(x) \neq 0\}$ must be a finite or countable subset of \mathbb{R} for each $\Psi \in \ell^2(\mathbb{R})$.

For every $x_0 \in \mathbb{R}$, let us denote by $\delta_{x_0} \in \ell^2(\mathbb{R})$ the characteristic function of the set $\{x_0\} \subset \mathbb{R}$, i.e.

$$\delta_{x_0}(x) := \begin{cases} 1 & \text{if } x = x_0 \\ 0 & \text{if } x \neq x_0. \end{cases}$$

The family $\{\delta_x : x \in \mathbb{R}\}$ constitutes an orthonormal basis of $\ell^2(\mathbb{R})$ and, hence, any $\Psi \in \ell^2(\mathbb{R})$ can be written in the form

$$\Psi = \sum_{x \in \mathbb{R}} \langle \delta_x, \Psi \rangle_{\ell^2(\mathbb{R})} \delta_x = \sum_{x \in \mathbb{R}} \Psi(x) \delta_x.$$

Let us denote by $\nu_{\mathfrak{b}\mathbb{R}}$ the probabilistic Haar measure on $\mathfrak{b}\mathbb{R}$ (the Bohr compactification of the real line) and consider the Hilbert space $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ with scalar product

$$\langle \psi_1, \psi_2 \rangle_{\mathfrak{b}\mathbb{R}} = \int_{\mathfrak{b}\mathbb{R}} \overline{\psi_1} \psi_2 \nu_{\mathfrak{b}\mathbb{R}}.$$

The space $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ is isomorphic to the Hilbert space $AP(\mathbb{R}, \mathbb{C})$ of the (complex) Besicovitch almost periodic functions on \mathbb{R} , with scalar product given by

$$\langle \psi_1, \psi_2 \rangle_{AP} = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T \overline{\psi_1(y)} \psi_2(y) dy.$$

The Fourier–Bohr transform

$$\mathfrak{B} : L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}}) \rightarrow \ell^2(\mathbb{R})$$

is an unitary isomorphism between $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ and $\ell^2(\mathbb{R})$. Explicitly, for every $\psi \in L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ we have

$$\mathfrak{B}(\psi)(x) := \langle \chi_x, \psi \rangle_{\mathfrak{b}\mathbb{R}} = \int_{\mathfrak{b}\mathbb{R}} \chi_{-x} \psi \nu_{\mathfrak{b}\mathbb{R}}$$

where, for every $x \in \mathbb{R}$, χ_x denotes the continuous character of $\mathfrak{b}\mathbb{R}$ such that $\chi_x(\iota(y)) = e^{ixy} =: \phi_x(y)$, for all $y \in \mathbb{R}$ and $\iota: \mathbb{R} \hookrightarrow \mathfrak{b}\mathbb{R}$ is the embedding of \mathbb{R} in $\mathfrak{b}\mathbb{R}$. The scalar product becomes now

$$\langle \psi_1, \psi_2 \rangle_{\mathfrak{b}\mathbb{R}} = \sum_{x \in \mathbb{R}} \overline{\mathfrak{B}(\psi_1)(x)} \mathfrak{B}(\psi_2)(x),$$

where it is important to notice that

$$\mathfrak{B}(\chi_{x_0}) = \delta_{x_0}, \quad x_0 \in \mathbb{R}$$

so that $\{\chi_x : x \in \mathbb{R}\}$ is an orthonormal basis of $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ and any $\psi \in L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ can be written in a unique form as

$$\psi = \sum_{x \in \mathbb{R}} \Psi(x) \chi_x, \quad \text{where } \Psi(x) = \mathfrak{B}(\psi)(x), \quad \Psi \in \ell^2(\mathbb{R}).$$

Equivalently, given $\psi \in AP(\mathbb{R}, \mathbb{C})$, its Fourier–Bohr transform

$$\mathfrak{B}(\psi)(x) := \langle \phi_x, \psi \rangle_{AP} = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T e^{-ixy} \psi(y) dy.$$

belongs to $\ell^2(\mathbb{R})$ and $\{\phi_x : x \in \mathbb{R}\}$ is an orthonormal basis of $AP(\mathbb{R}, \mathbb{C})$.

3. Polymer representations for the Weyl algebra

The Weyl algebra, codified by the equation

$$U(p)V(q) = e^{-ipq}V(q)U(p), \quad \forall q, p \in \mathbb{R},$$

admits infinitely many unitarily inequivalent irreducible representations (see, for example [15]). In this paper we will use two unitarily inequivalent irreducible representations on $\ell^2(\mathbb{R})$ that we refer to as position and momentum representations, respectively [11]:

PR of the Weyl algebra on $\ell^2(\mathbb{R})$. The one-parameter families of unitary operators $\{U(p) : p \in \mathbb{R}\}$ and $\{V(q) : q \in \mathbb{R}\}$ defined through

$$U(p)\delta_{q_0} := e^{ipq_0}\delta_{q_0}, \quad V(q)\delta_{q_0} := \delta_{q_0-q}, \quad \forall p, q, q_0 \in \mathbb{R},$$

satisfy

$$U(p)V(q) = e^{-ipq}V(q)U(p), \quad \forall q, p \in \mathbb{R}.$$

In this representation, $q \mapsto V(q)$ is not weakly continuous and there is no momentum operator. On the other hand, the map $p \mapsto U(p)$ is weakly continuous and there is a position self-adjoint unbounded operator $Q := -iU'(0)$, satisfying

$$Q\delta_q = q\delta_q,$$

with domain

$$\mathcal{D}(Q) := \left\{ \Psi \in \ell^2(\mathbb{R}) : \sum_{q \in \mathbb{R}} q^2 |\Psi(q)|^2 < \infty \right\}.$$

MR of the Weyl algebra on $\ell^2(\mathbb{R})$. The one-parameter families of unitary operators $\{U(p) : p \in \mathbb{R}\}$ and $\{V(q) : q \in \mathbb{R}\}$ defined through

$$U(p)\delta_{p_0} := \delta_{p_0+p}, \quad V(q)\delta_{p_0} := e^{iqp_0}\delta_{p_0}, \quad \forall q, p, p_0 \in \mathbb{R},$$

satisfy

$$U(p)V(q) = e^{-ipq}V(q)U(p), \quad \forall q, p \in \mathbb{R}.$$

In this representation, $p \mapsto U(p)$ is not weakly continuous and there is no position operator. On the other hand, the map $q \mapsto V(q)$ is weakly continuous and there is a momentum self-adjoint unbounded operator $\mathbf{P} := -iV'(0)$, satisfying

$$\mathbf{P}\delta_p = p\delta_p,$$

with domain

$$\mathcal{D}(\mathbf{P}) := \left\{ \Psi \in \ell^2(\mathbb{R}) : \sum_{p \in \mathbb{R}} p^2 |\Psi(p)|^2 < \infty \right\}.$$

By using the Fourier–Bohr transform, both representations can be implemented in $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$. In particular, if we use $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ as the carrier of the PR, the operator \mathbf{Q} is just a derivative operator and $\mathbf{Q}^2 = -\mathbf{\Delta}$, where $\mathbf{\Delta}$ is the Laplace operator. On the other hand, in the MR, the operator \mathbf{P} is just a derivative operator and $\mathbf{P}^2 = -\mathbf{\Delta}$.

As we will see later when we study the spectra of the polymerized harmonic oscillator Hamiltonians, the eigenvalue equations can be equivalently written as *difference equations* in $\ell^2(\mathbb{R})$ or as *differential equations* in $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ with no reason to prefer, *a priori*, one writing over the other. This is in exact analogy with the relationship between the matrix mechanics of Heisenberg and the wave mechanics of Schrödinger.

4. Polymerized Hamiltonians: general results on the spectra

Let $W : \mathbb{R} \rightarrow \mathbb{R}$ be a periodic piecewise continuous (in the standard topology for \mathbb{R}) function, then W defines, through the multiplication in the algebra of almost periodic functions, a bounded operator \mathbf{W} in $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ (equivalently in $AP(\mathbb{R}, \mathbb{C})$). The operator

$$\mathbf{H} = -\mathbf{\Delta} + \mathbf{W} \tag{1}$$

is a self-adjoint unbounded operator with domain

$$\mathcal{D}(\mathbf{H}) = \mathcal{D}(\mathbf{\Delta}) = \left\{ \psi \in L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}}) : \sum_{x \in \mathbb{R}} x^4 |\mathfrak{B}(\psi)(x)|^2 < \infty \right\}.$$

The crucial result that allows us to obtain the spectrum of (1) in a straightforward way can be found in [16–19] where it is shown that the operator (1) has only *pure point spectrum* that coincides with the spectrum of the Schrödinger operator $H = -\Delta + W$, with periodic potential W , defined on $L^2(\mathbb{R}, \mu_{\mathbb{R}})$. We can then use the following well-known theorem [13] (that encapsulates the Bloch theorem of the physicists or the Floquet theory of the mathematicians):

Theorem. *Let W be a piecewise continuous function of period 2π . Let $H = -\Delta + W$ on $L^2(\mathbb{R}, \mu_{\mathbb{R}})$. Let $E_1(0), E_2(0), \dots$ be the eigenvalues of the corresponding operator on $(0, 2\pi)$ with periodic boundary conditions and let $E_1(\pi), E_2(\pi), \dots$ be the eigenvalues with antiperiodic boundary conditions. Let*

$$E_n^L = \begin{cases} E_n(0) & n \text{ odd} \\ E_n(\pi) & n \text{ even} \end{cases} \quad E_n^R = \begin{cases} E_n(\pi) & n \text{ odd} \\ E_n(0) & n \text{ even} \end{cases}.$$

Then H has purely absolutely continuous spectrum and

$$\sigma(H) = \sigma_{\text{ac}}(H) = \bigcup_{n=1}^{\infty} [E_n^L, E_n^R].$$

Hence we conclude that

$$\sigma(\mathbf{H}) = \sigma_{\text{pp}}(\mathbf{H}) = \sigma(H) = \sigma_{\text{ac}}(H) = \bigcup_{n=1}^{\infty} [E_n^L, E_n^R]. \quad (2)$$

Moreover, the eigenvalues of \mathbf{H} are, at most, doubly degenerate [18]. Associated with the bands (2) in the energy spectrum there are spectral gaps of forbidden energies

$$(-\infty, E_1^L) \cup (E_1^R, E_2^L) \cup (E_2^R, E_3^L) \cup \dots$$

Notice that some of the gaps will be absent if the edges E_n^R and E_{n+1}^L of two consecutive bands coincide. This characterization of the spectrum as consisting of a set of bands is the main result of the paper. We would like to mention here that this fact is somehow implicit in [1] (see section 3.2.) although, to our knowledge, the structure of the spectrum of the Hamiltonian as explained above has not been made explicit in the literature.

It is not difficult to understand the previous result from an intuitive point of view. Indeed, the almost periodic solutions that are found by solving the eigenvalue equation (in its differential form) are *not normalizable* in the standard $L^2(\mathbb{R}, \mu_{\mathbb{R}})$ sense but *they are* when the scalar product in the space of almost periodic functions is used. Although it is true that strictly periodic (or anti-periodic) solutions solve the preceding equation there is a continuum of solutions parametrized by a real number—this is the usual statement that underlies the Bloch–Floquet theorem. Indeed, this theorem shows that the non-normalizable ‘eigenfunctions’ φ of $H = -\Delta + W$ satisfy

$$\varphi(x + 2\pi) = e^{2\pi k i} \varphi(x), \quad \forall x \in \mathbb{R}, \quad (3)$$

for some real number k —that can be restricted to lie in $k \in [0, 1)$ or, equivalently, in $k \in [-1/2, 1/2)$. The choice $k = 0$ leads to the $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ eigenvectors corresponding to ‘half’ of the boundary points of the band spectrum (2π -periodic solutions, that determine either E_{2n+1}^L or E_{2n}^R). The choice $k = 1/2$ leads to the remaining boundary points (2π -antiperiodic solutions, that determine either E_{2n}^L or E_{2n+1}^R). The other choices for k give rise to the eigenfunctions corresponding to the rest of the points in the energy bands.

It is straightforward to show that if φ satisfies (3) it is possible to write

$$\varphi(x) = e^{ikx} f(x)$$

for some 2π -periodic function f . Then, it is obvious that its Fourier–Bohr transform

$$\mathfrak{B}(\varphi)(x) = \lim_{N \rightarrow \infty} \frac{1}{2\pi N} \int_{-N\pi}^{N\pi} e^{-ixy} \varphi(y) dy = \lim_{N \rightarrow \infty} \frac{1}{2\pi N} \int_{-N\pi}^{N\pi} e^{-ixy} e^{iky} f(y) dy$$

vanishes when

$$x \not\equiv k \pmod{1}.$$

In other words, if $\varphi = \varphi_k$ satisfies (3) for some k , the set $\{x \in \mathbb{R} : \mathfrak{B}(\varphi_k)(x) \neq 0\}$ is a subset of the regular lattice $\gamma_k = \{k + n : n \in \mathbb{Z}\} \subset \mathbb{R}$. As we will see in the next section, this fact has a neat consequence when the eigenvalue equation for the polymer Hamiltonian \mathbf{H} is written in $\ell^2(\mathbb{R})$: as elements of $\ell^2(\mathbb{R})$ the eigenfunctions have their support on *regular, evenly spaced*, lattices of the previous form. If $k_1 \not\equiv k_2 \pmod{1}$ the lattices γ_{k_1} and γ_{k_2} are disjoint and, hence, the states φ_{k_1} and φ_{k_2} are orthogonal. As a final comment we would like to point out here that it is simply not true that the eigenvalue problem for a polymer Hamiltonian $\mathbf{H} = -\Delta + W$ is equivalent to the one corresponding to a particle on a unit circle, whose solutions satisfy $\varphi(x + 2\pi) = e^{2\pi k i} \varphi(x)$ only for $k = 0$.

5. Different representations for the polymerized harmonic oscillator

The Hamiltonian for the classical harmonic oscillator, written in terms of dimensionless variables q and p , has the form

$$H(q, p) = \frac{\hbar^2}{2m\ell^2} p^2 + \frac{m\ell^2\omega^2}{2} q^2,$$

where ℓ has dimension of length, m has dimension of mass, ω is the frequency of the harmonic oscillator and \hbar is the reduced Planck constant. The main difference between the quantizations for the harmonic oscillators in the standard Schrödinger representations of the Weyl algebra and the polymer PR or MR is the fact that the Hamiltonian in the latter cases cannot be represented because it is not possible to simultaneously quantize the q^2 and the p^2 parts of the Hamiltonian. In order to build the quantum Hamiltonian operator let us work in the PR. The first step [1] (see also [12]) is to introduce an arbitrary, but fixed, parameter $q_0 > 0$ (that defines a length scale $q_0\ell$) and consider the following one-parameter family of operators:

$$\mathbf{H}(q_0) := \frac{\hbar^2}{2m(2q_0\ell)^2} (2 - \mathbf{V}(2q_0) - \mathbf{V}(-2q_0)) + \frac{m\ell^2\omega^2}{2} \mathbf{Q}^2. \quad (4)$$

By defining the self-adjoint operators

$$\mathbf{P}(q_0) := \frac{i}{2q_0} (\mathbf{V}(q_0) - \mathbf{V}(-q_0))$$

these Hamiltonians can be written as

$$\mathbf{H}(q_0) := \frac{\hbar^2}{2m\ell^2} \mathbf{P}(q_0)^2 + \frac{m\ell^2\omega^2}{2} \mathbf{Q}^2$$

and, hence, it is straightforward to see that for every value of $q_0 > 0$ these are non-negative operators in $\ell^2(\mathbb{R})$ with the same domain as $-\mathbf{\Delta}$.

Alternatively, in the MR, the quantum harmonic oscillator Hamiltonian can be modeled by

$$\mathbf{H}(p_0) := \frac{\hbar^2}{2m\ell^2} \mathbf{P}^2 + \frac{m\ell^2\omega^2}{2(2p_0)^2} (2 - \mathbf{U}(2p_0) - \mathbf{U}(-2p_0)), \quad (5)$$

where $p_0 > 0$ introduces a momentum scale $p_0\hbar/\ell$. If the momentum and length scales defined by the parameter q_0 and p_0 satisfy

$$\frac{\hbar}{\ell} p_0 = m\omega q_0\ell$$

the spectra of (4) and (5) coincide so we will use (4) in the following. (Notice, however, that the MR is the one used in many polymer quantum field theories [6–8]). A comment is in order here: the approximation that is used in (4) to make sense of p^2 in the polymer PR can be seen as just a simple and natural choice among a large family of possible ones (as mentioned in [12]); another such—and natural—choice will be considered below.

We are interested in studying the spectrum of the Hamiltonian (4) as a function of the parameter q_0 . To this end we will write the eigenvalue equation both in the $\ell^2(\mathbb{R})$ and $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ PRs. The eigenvalue equation in the $\ell^2(\mathbb{R})$ PR is

$$\frac{\hbar^2}{2m(2q_0\ell)^2} (2\Psi(q) - \Psi(q + 2q_0) - \Psi(q - 2q_0)) + \frac{m\ell^2\omega^2}{2} q^2\Psi(q) = E\Psi(q), \quad \forall q \in \mathbb{R}$$

where the solutions $\Psi \neq 0$ must satisfy the normalizability condition:

$$\sum_{q \in \mathbb{R}} |\Psi(q)|^2 < \infty, \quad \sum_{q \in \mathbb{R}} q^4 |\Psi(q)|^2 < \infty,$$

and the non-negativity of the Hamiltonian guarantees that the only possible eigenvalues E must satisfy $E \geq 0$. This is a difference equation that establishes a relationship between the values of Ψ at three different points q , $q + 2q_0$ and $q - 2q_0$. Actually, as mentioned at the end of section 4 (see also [1]), it is possible to solve this difference equation by looking for solutions Ψ_k , where $k \in [0, 1)$, for which $\{q \in \mathbb{R} : \Psi_k(q) \neq 0\}$ is a subset of the regular lattice

$$2q_0 \cdot \gamma_k := \{2q_0k + 2q_0n : n \in \mathbb{Z}\}.$$

On the other hand, the eigenvalue equation in $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ is the differential equation

$$\psi''(p) + \left(\frac{2E}{m\ell^2\omega^2} - \frac{\hbar^2}{m^2\ell^4\omega^2q_0^2} \sin^2(q_0p) \right) \psi(p) = 0 \quad (6)$$

that must be solved for those $\psi \in L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ belonging to the domain of the operator $\mathcal{Q}^2 = -\Delta$. By performing the change of variables

$$x = 2q_0p$$

and defining $\varphi(x)$ such that

$$\psi(p) = \varphi(2q_0p), \quad \psi''(p) = 4q_0^2\varphi''(2q_0p),$$

the eigenvalue equation (6) can be rewritten in the form of the familiar Mathieu equation

$$\varphi''(x) + \left(\frac{E}{2m\omega^2(q_0\ell)^2} - \frac{1}{2} \left(\frac{\hbar}{2m\omega(q_0\ell)^2} \right)^2 (1 - \cos x) \right) \varphi(x) = 0 \quad (7)$$

for $\varphi \in L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$.

Notice that both equations (6) and (7) can be rewritten as an eigenvalue problem of the form

$$(-\Delta + \mathbf{W}(q_0))\psi = \lambda\psi, \quad \psi \in L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}}).$$

In particular, for equation (6) we have

$$\lambda = \frac{2E}{m\ell^2\omega^2}$$

and the bounded operators $\mathbf{W}(q_0)$ are defined by the (π/q_0) -periodic functions $W(\cdot|q_0) : \mathbb{R} \rightarrow \mathbb{R}$:

$$W(p|q_0) = \frac{\hbar^2}{m^2\ell^4\omega^2q_0^2} \sin^2(q_0p).$$

On the other hand, equation (7) is written in terms of the 2π -periodic potentials $w(\cdot|q_0) : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$w(x|q_0) := \frac{1}{2} \left(\frac{\hbar}{2m\omega(q_0\ell)^2} \right)^2 (1 - \cos x).$$

It is very important to emphasize at this point that, as stated in section 4, in order to obtain the energy spectrum, we must find the band spectrum of the corresponding Schrödinger operators in $L^2(\mathbb{R}, \mu_{\mathbb{R}})$. We would like to mention here that the equation (7), considered as an eigenvalue problem in the separable Hilbert space $L^2(\mathbb{S}^1, \mu_{\mathbb{S}^1})$, where $\mu_{\mathbb{S}^1}$ is the Haar measure for \mathbb{S}^1 , was proposed in [20] to study the quantum Harmonic oscillator on a lattice. Notice that the non-trivial solutions φ to equation (7) that satisfy $\varphi \in L^2(0, 2\pi)$, $\varphi(2\pi) = \varphi(0)$, and $\varphi'(2\pi) = \varphi'(0)$ are just the solutions to (7) in $L^2(\mathbb{S}^1, \mu_{\mathbb{S}^1})$. These are also relevant when (7) is considered in $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ because they provide the eigenvectors corresponding to certain boundary points of the band spectrum (either E_{2n+1}^L or E_{2n}^R). The antiperiodic solutions

$\varphi \in L^2(0, 2\pi)$, satisfying $\varphi(2\pi) = -\varphi(0)$ and $\varphi'(2\pi) = -\varphi'(0)$, provide the eigenvectors corresponding to the remaining boundary points of the bands (either E_{2n}^L or E_{2n+1}^R). It is important to notice that, in addition to these, there are solutions $\varphi \in L^2(0, 2\pi)$ satisfying $\varphi(2\pi) = e^{2\pi ki}\varphi(0)$ and $\varphi'(2\pi) = e^{2\pi ki}\varphi'(0)$, for any $k \in \mathbb{R}$, that give rise to the eigenfunctions of the rest of the points in energy bands.

The general results of section 4 guarantee that

$$\sigma(\mathbf{H}(q_0)) = \sigma_{\text{pp}}(\mathbf{H}(q_0)) = \bigcup_{n=1}^{\infty} [E_n^L(q_0), E_n^R(q_0)], \quad E_n^L(q_0) < E_n^R(q_0).$$

In this case, it is also known [21] that (see, also [1, 10])

$$\lim_{q_0 \rightarrow 0} E_n^R(q_0) = \lim_{q_0 \rightarrow 0} E_n^L(q_0) = E_n,$$

where E_n is the n th eigenvalue of the $L^2(\mathbb{R}, \mu_{\mathbb{R}})$ harmonic oscillator Hamiltonian

$$H_{\text{HO}} = -\frac{m\ell^2\omega^2}{2}\Delta + W_{\text{HO}},$$

where $W_{\text{HO}} : \mathbb{R} \rightarrow \mathbb{R}$ is the quadratic potential

$$W_{\text{HO}}(p) = \frac{\hbar^2}{m^2\ell^4\omega^2}p^2.$$

Notice, however, that for a fixed value of $q_0 > 0$ all the gaps between bands are present (see, for example, [13]), i.e.

$$D_n(q_0) := E_{n+1}^L(q_0) - E_n^R(q_0) > 0, \quad \forall n = 1, 2, \dots$$

Actually, the width of the gaps behaves as [22]

$$D_n(q_0) = \frac{m\omega^2(q_0\ell)^2}{4^{n-1}(n-1)!^2} \left(\frac{\hbar}{2m\omega(q_0\ell)^2} \right)^{2n} (1 + o(1/n^2)).$$

As we can see they get narrower and narrower as the energy grows. In practice this means that for high energies the spectrum basically consists of a continuum of eigenvalues belonging to the point spectrum (i.e. associated with *normalizable* states). This behavior is very different from that of the ordinary quantum harmonic oscillator.

A comment is in order now. Although the simplest way to model the standard harmonic oscillator Hamiltonian in the polymer PR is by approximating p^2 as

$$\mathbf{P}(q_0)^2 = \frac{m^2\ell^4\omega^2}{\hbar}\mathbf{W}(q_0), \quad q_0 > 0,$$

there is, in fact, a huge ambiguity here (as has been realized by other authors [12]). In particular, in the PR supported by $L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$, one can actually use, for example, any (π/q_0) -periodic function of $W(\cdot|q_0) : \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\lim_{q_0 \rightarrow 0} W(p|q_0) = \frac{\hbar^2}{m^2\ell^4\omega^2}p^2 \quad (\text{pointwise}).$$

A very simple such choice is provided by the (π/q_0) -periodic extension of a purely quadratic (pq) potential

$$W_{\text{pq}}(p|q_0) := \frac{\hbar^2}{m^2\ell^4\omega^2}p^2, \quad p \in \left[-\frac{\pi}{2q_0}, \frac{\pi}{2q_0} \right].$$

The band structure in this case is qualitatively equal to the one described for the Mathieu equation. As can be seen in figure 1, the first bands closely correspond to the usual eigenvalues of the harmonic oscillator and are very narrow. On the other hand it can be readily seen that as




Figure 1. Band structure for $H = -\Delta + W_{\text{pq}}$ where W_{pq} is the periodic potential satisfying $W_{\text{pq}}(x) = x^2$ for $x \in [-4, 4]$. This corresponds to choosing $q_0 = \pi/8$ and $\ell = \hbar = m = \omega = 1$. We have plotted the potential in one period, the bands that constitute the spectrum $\sigma(H)$, and the trace of the matrix $M(E)$ that determines the position of the bands as those values of the energy for which $\text{Tr } M(E) = \pm 2$ (see [13]). The narrow lowest energy bands closely correspond to the lowest quantum harmonic oscillator eigenvalues.

the energy grows the bands widen and the gaps become more and more narrow, in agreement with the behavior described above in the case of the Mathieu approximation.

An interesting question that arises at this point is whether all these possible extensions behave as the standard quantum harmonic oscillator in the $q_0 \rightarrow 0$ limit or, more precisely, how do all the different spectra compare in this limit. This question is made even more relevant by the fact that there are periodic potentials (leading, for example, to the so-called Lamé equation) that give rise only to a finite number of bands in the spectrum. If such potentials could be used to approximate a periodic extension of a purely quadratic one with arbitrary large period it is difficult to understand to what extent the spectrum of the ordinary quantum harmonic oscillator could be recovered. A partial answer can be given by considering pairs of Hamiltonians that are close to each other and using perturbation theory. If one takes $\mathbf{H}_0 = -\Delta + \mathbf{W}_0, \mathbf{H}_1 = -\Delta + \mathbf{W}_1$ where $\mathbf{W}_1 = \mathbf{W}_0 + \mathbf{V}$ and $\|\mathbf{V}\| < \epsilon$ then $\mathbf{H}_1 = \mathbf{H}_0 + \mathbf{V}$ and the eigenvalues of \mathbf{H}_1 are approximated by those of \mathbf{H}_0 with $O(\epsilon)$ corrections. Notice, however, that if there is no need to recover p^2 for all $p \in \mathbb{R}$, *any potential* term that has a quadratic minimum at $p = 0$ of the same type as the one in the potential that appears in the Mathieu equation provides a perfectly valid approximation for the harmonic oscillator Hamiltonian at small energies. An interesting example in this respect is provided by periodic potentials defined by Jacobian elliptic functions, in particular those leading to the Lamé equation [23]. It is known that, in those cases the spectrum consists of a continuum of positive eigenvalues with a finite number of gaps (or, equivalently, on a finite number of bounded bands and an unbounded band). These examples show that one has to be careful when extrapolating the form of the spectrum from the band structure of the polymerized quantum Hamiltonians for the harmonic oscillator.

6. Statistical mechanics of the polymerized harmonic oscillator

We briefly discuss in this section the statistical mechanics of a single particle whose quantum dynamics is governed by a Hamiltonian $\mathbf{H} = -\Delta + \mathbf{W}$ belonging to the class (1). We will show that neither the microcanonical nor the canonical ensembles provide a convenient starting point to study the statistical mechanics of the system. This is so because both the statistical (counting) entropy and the partition function for the system are ill-defined. Notice that, at face value, this would imply that fundamental model systems such as the Einstein crystal would be ill-defined if the polymer quantization of the harmonic oscillator is used.

For $E \in \sigma(\mathbf{H}) = \sigma_{\text{pp}}(\mathbf{H})$, let us denote by $\mathbf{P}_{\mathbf{H}}(E)$ the self-adjoint orthogonal projector on the eigenspace $\mathbf{H} = E$. These projectors give rise to the resolution of the identity

$$\mathbf{I} = \sum_{E \in \sigma(\mathbf{H})} \mathbf{P}_{\mathbf{H}}(E) \quad \text{i.e.} \quad \psi = \sum_{E \in \sigma(\mathbf{H})} \mathbf{P}_{\mathbf{H}}(E)\psi, \quad \forall \psi \in L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}}).$$

The statistical entropy for the system as a function of the energy is given by

$$\Omega(E) := \text{Tr} \left(\sum_{\epsilon \in \sigma(\mathbf{H}) \cap (-\infty, E]} \mathbf{P}_{\mathbf{H}}(\epsilon) \right) = \sum_{\epsilon \in \sigma(\mathbf{H}) \cap (-\infty, E]} d(\epsilon)$$

where $d(E)$ is the degeneracy of the energy eigenvalue E . As we mentioned above (see [18]) $d(E)$ is at most two for generic one-dimensional problems described by (1). As it is plainly obvious the continuous sums written above are ill-defined owing to the uncountable number of energy eigenstates of the system. Likewise the canonical partition function is also ill-defined. In this case we have to consider (by using the spectral representation theorem)

$$e^{-\beta \mathbf{H}} = \sum_{E \in \sigma(\mathbf{H})} e^{-\beta E} \mathbf{P}_{\mathbf{H}}(E), \quad \beta > 0.$$

Notice that $e^{-\beta H}$ is a bounded operator because

$$\|e^{-\beta H} \psi\|^2 = \sum_{E \in \sigma(\mathbf{H})} e^{-2\beta E} \|\mathbf{P}_H(E) \psi\|^2 \leq \sum_{E \in \sigma(\mathbf{H})} \|\mathbf{P}_H(E) \psi\|^2 = \|\psi\|^2, \quad \forall \psi \in L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$$

but it is not a trace class operator,

$$Z(\beta) = \text{Tr}(e^{-\beta H}) = \sum_{E \in \sigma(\mathbf{H})} d(E) e^{-\beta E} = \infty,$$

because $\#\{E \in \sigma(\mathbf{H}) : d(E)e^{-\beta E} > 0\} = \#\mathbb{R}$. Hence the thermal density matrix

$$\rho = \frac{e^{-\beta H}}{\text{Tr}(e^{-\beta H})}$$

is ill-defined.

Notice that the situation that we are facing here is conceptually different from the one corresponding to a standard Schrödinger operator $H = -\Delta + W$ in $L^2(\mathbb{R}, \mu_{\mathbb{R}})$ for a one-dimensional periodic potential W . Indeed, despite the fact that in that case, as discussed in section 4, the spectrum of H is *purely absolutely continuous*, bounded from below, and the operators $e^{-\beta H}$ are trace class for $\beta > 0$. In particular, the microcanonical entropy is given by

$$\Omega(E) = \int_{\sigma(H) \cap (-\infty, E]} g(\epsilon) d\epsilon$$

in terms of $g(E)$, the density of states per energy interval, and the partition function is

$$Z(\beta) = \int_{\sigma(H)} g(E) e^{-\beta E} dE.$$

The statistical entropy is obviously well defined, the partition function too (as long as the density of states $g(E)$ does not grow too fast) and we have the standard probabilistic interpretation associated with the canonical ensemble:

$$\text{Prob}(B|\beta) = \frac{1}{Z(\beta)} \int_B g(E) e^{-\beta E} dE, \quad B \subset \sigma(H).$$

A possible way out of the situation explained above would be to find a physical criterion to eliminate most of the eigenstates of the Hamiltonian and just leave a countable number of them. Suppose that, for every $n \in \mathbb{N}$, we select one energy eigenvalue in each band, $E_n \in [E_n^L, E_n^R]$, with $E_n < E_{n+1}$ and $\lim_{n \rightarrow \infty} E_n = \infty$. In general, let

$$\mathcal{E} := \bigcup_{n=1}^{\infty} \{E_n\}, \quad \mathbf{P}_{\mathcal{E}} := \sum_{E \in \mathcal{E}} \mathbf{P}_H(E), \quad \mathbf{H}_{\mathcal{E}} := \mathbf{H} \mathbf{P}_{\mathcal{E}} = \mathbf{P}_{\mathcal{E}} \mathbf{H} \mathbf{P}_{\mathcal{E}},$$

then

$$\Omega_{\mathcal{E}}(E) := \text{Tr} \left(\sum_{\epsilon \in \mathcal{E} \cap (-\infty, E]} \mathbf{P}_H(\epsilon) \right) = \sum_{\epsilon \in \mathcal{E} \cap (-\infty, E]} d(\epsilon) = \sum_{E_n \leq E} d(E_n)$$

and

$$Z_{\mathcal{E}}(\beta) = \text{Tr}(e^{-\beta \mathbf{H}_{\mathcal{E}}}) = \sum_{E \in \mathcal{E}} d(E) e^{-\beta E} = \sum_{n=1}^{\infty} d(E_n) e^{-\beta E_n} < \infty, \quad \forall \beta > 0,$$

are well defined.

We would like to mention here that, within the context of LQG, it is customary to talk about superselected sectors in the full gravity Hilbert space. A mechanism that may be at work in this setting is the selection of a separable physical subspace of the full non-separable Hilbert space in the process of implementing the quantum constraints *à la Dirac*. This has been

checked in detail in the context of the polymer quantization of parameterized field theories by Laddha and Varadarajan [25, 26].

Superselection rules may be the ingredient needed to avoid some of the statistical problems mentioned above. In particular, given $k \in [0, 1)$, let us consider [1] the subspace $\mathcal{H}_k \subset L^2(\mathfrak{b}\mathbb{R}, \nu_{\mathfrak{b}\mathbb{R}})$ that is generated by the energy eigenfunctions satisfying $\varphi(x + 2\pi) = e^{2\pi ki}\varphi(x)$. The Fourier–Bohr transform of any of such eigenvectors fulfils

$$\{x \in \mathbb{R} : \mathfrak{B}(\varphi)(x) \neq 0\} \subset \gamma_k = \{k + n : n \in \mathbb{Z}\}$$

and it is possible to prove that each subspace \mathcal{H}_k is ‘superselected’ in the sense that it is not only trivially invariant under the action of \mathbf{H} but it is also invariant under both the action of the corresponding self-adjoint position operator and that of an appropriately chosen self-adjoint operator that approximates the momentum in the polymer PR [1]. (However, it is important to notice that it is possible to approximate the momentum in such a way that the subspaces superselected in that way become $\mathcal{H}_k \oplus \mathcal{H}_{1/2-k}$, for $k \in [0, 1/2)$, or even more general combinations.) The canonical ensemble defined by the restriction of the harmonic oscillator Hamiltonian to one of such \mathcal{H}_k -sectors has been considered in [20]. However, notice that, as mentioned in standard references on superselection rules (see, for instance, [24]), it is perfectly acceptable to have density matrices involving states in different superselected sectors so, in principle, one should not exclude statistical ensembles including such states. The existence of superselected subspaces in a physical Hilbert space, rather than precluding the possibility of considering non-trivial linear combinations of states in different sectors, entails the *physical impossibility* of measuring the relative phases between them. This, in turn, means that such states are physically realizable only as density matrices. In any case, unlike the situation in LQG, in the simple quantum mechanical model that we have considered in the paper, this recourse to superselection rules does not seem to be justified. This is so because, if its sole purpose is to render the Hilbert space separable by restricting the quantum dynamics to a Hilbert subspace consisting of an orthonormal sum of eigenspaces of the Hamiltonian, there is no point in introducing a polymer quantization to begin with. Standard quantum mechanics would suffice.

7. Final comments and conclusions

The main result discussed in the paper is the fact that, generically, the spectrum of the polymer Hamiltonians for the harmonic oscillator consists of *bands* similar to the ones appearing in the study of periodic potentials in ordinary quantum mechanics. This means that the limit in which the length scale defined by q_0 becomes very small should be handled with proper care. Although there is evidence supporting that the energy eigenvalues for the standard quantum harmonic oscillator are recovered in this limit, important features of the spectrum differ in significant ways, the most important one being the fact that the effective degeneracy of the energy eigenvalues becomes infinite (an *uncountable infinite*, in fact).

The underlying reason for this is easy to understand: the differential symbol for the eigenvalue equation in the PR is the standard one for a particle in a periodic potential. The fact that the scalar product is not the standard one renders the solutions to the equation (for suitable values of the energy) normalizable (whereas they are not if the standard scalar product in $L^2(\mathbb{R}, \mu_{\mathbb{R}})$ is used). In this sense the periodic and antiperiodic solutions that are mentioned in the literature are just a small subset of all the (almost-periodic) solutions permitted by the Bloch theorem. Notice, by the way, that it is not correct to think of the polymerized Hamiltonians as corresponding to a particle in a circle with a potential with a second order minimum (where only periodic—and not antiperiodic—‘boundary conditions’ would make sense). The correct

point of view is to think about such systems as consisting of a particle in a periodic potential with wave functions defined in the Hilbert space of almost periodic functions.

There are a number of results in the literature regarding the band structure for periodic potentials [13] and proofs of specific properties of the spectrum for particular choices. The details of how the limit in which the parameter q_0 goes to zero works are subtle. As has been shown in the examples of section 5 the number of bands appearing in the spectrum for one-dimensional polymer Hamiltonians of the type that we have discussed here may be finite (as in the case of the Lamé potentials, see [23] and references therein) or infinite as it happens in the case of the usual Hamiltonian leading to the Mathieu equation.

The quantization of many free theories relies heavily on the quantum harmonic oscillator as these systems can be thought of as consisting of an infinite number of such systems. A possible approach to the quantization of these models [10] is based on the construction of a Fock space on the polymer Hilbert space considered here (see [6–9] for different approaches). In principle it is to be expected that the band structure that we have discussed may play an important role and have observable consequences. The easiest way to explore those is by looking to two-point functions and, very especially, to field commutators. It is interesting, in particular, to understand if the polymerization changes the microcausality of the system.

The impossibility of defining a partition function for the polymerized models discussed here by generalizing the standard expressions in quantum statistical mechanics (involving traces and trace-class operators) is, to say the least, disturbing and we do not see an obvious way out in this setting. Notice, however, that this negative conclusion should not be naively carried over to LQG. First of all, as mentioned above, the final physical Hilbert space of LQG can very well be separable despite the fact that the kinematical Hilbert space is not thus avoiding any difficulties of the type explained above. Also, at least from the point of view of gravity, the fact that one cannot define the energy in a natural way leads naturally to the consideration of ensembles associated with operators different from the Hamiltonian, in particular the area operator (with its discrete spectrum and effectively finite degeneracies) that is customarily used, for example, to discuss black hole entropy in LQG [27–29]. This change of ensemble may be what is needed to have a sensible framework for the statistical mechanics of quantum gravitational systems [30, 31].

To end we would like to mention that, although it is not straightforward to extend the results of the present paper to LQC (and, actually, they may not be directly applicable) we think that it may be instructive to look at the differential version of the difference equations that are usually employed to discuss the quantum dynamics of the early universe in LQC. In that context the argued presence of superselection rules (that remove huge sectors from the space of quantum states) seems to sidestep some of the difficulties that crop up in the simple model provided by the polymerized harmonic oscillator. We think, however, that the issues that we have raised in the paper may be relevant for more advanced models incorporating more degrees of freedom.

Acknowledgments

We would like to thank W Chojnacki, A Corichi, J Lewandowski, T Pawłowski, P Singh and M Varadarajan for their valuable comments. This work has been supported by the Spanish MICINN research grants FIS2009-11893, FIS2012-34379 and the Consolider-Ingenio 2010 Program CPAN (CSD2007-00042).

References

- [1] Ashtekar A, Fairhurst S and Willis J L 2003 Quantum gravity, shadow states, and quantum mechanics *Class. Quantum Grav.* **20** 1031–62 (arXiv:gr-qc/0207106)
- [2] Corichi A, Vukašinac T and Zapata J A 2007 Hamiltonian and physical Hilbert space in polymer quantum mechanics *Class. Quantum Grav.* **24** 1495–512 (arXiv:gr-qc/0610072)
- [3] Corichi A, Vukašinac T and Zapata J A 2007 Polymer quantum mechanics and its continuum limit *Phys. Rev. D* **76** 044016 (arXiv:0704.0007 [gr-qc])
- [4] Ashtekar A and Singh P 2011 Loop quantum cosmology: a status report *Class. Quantum Grav.* **28** 213001 (arXiv:1108.0893)
- [5] Bojowald M 2008 Loop quantum cosmology *Living Rev. Rel.* **11** 4
- [6] Ashtekar A, Lewandowski J and Sahlmann H 2003 Polymer and Fock representations for a scalar field *Class. Quantum Grav.* **20** L11–1 (arXiv:gr-qc/0211012)
- [7] Kamiński W, Lewandowski J and Bobieński M 2006 Background independent quantizations: the scalar field I *Class. Quantum Grav.* **23** 2761–70 (arXiv:gr-qc/0508091)
- [8] Kamiński W, Lewandowski J and Okołów A 2006 Background independent quantizations: the scalar field II *Class. Quantum Grav.* **23** 5547–86 (arXiv:gr-qc/0604112)
- [9] Domagała M, Dziendzikowski M and Lewandowski J 2011 *Proceedings of the 3rd Quantum Geometry and Quantum Gravity School in Zakopane* arXiv:1210.0849v1 [gr-qc]
- [10] Hossain G M, Husain V and Seahra S S 2010 Propagator in polymer quantum field theory *Phys. Rev. D* **82** 124032 (arXiv:1007.5500 [gr-qc])
- [11] Halvorson H 2004 Complementarity of representations in quantum mechanics *Stud. Hist. Phil. Mod. Phys.* **35** 45–56 (arXiv:quant-ph/0110102)
- [12] Corichi A and Vukašinac T 2012 Effective constrained polymeric theories and their continuum limit arXiv:1202.1846v2 [gr-qc]
- [13] Reed M and Simon B 1978 *Methods of Modern Mathematical Physics* vol 4 (New York: Academic)
- [14] Chacón-Acosta G, Manrique E, Dagdug L and Morales-Técolt H A 2011 Statistical thermodynamics of polymer quantum systems *SIGMA* **7** 110–33 (arXiv:1109.0803v2 [gr-qc])
- [15] Chojnacki W 1987 Spectral analysis of Schrödinger operators in non-separable Hilbert spaces *Rend. Circ. Mat. Palermo (2) Suppl.* **17** 135–51
- [16] Chojnacki W 1986 Almost periodic Schrödinger operators in $L^2(b\mathbb{R})$ whose point spectrum is not all of the spectrum *J. Funct. Anal.* **65** 236–42
- [17] Burnat M 1964 Die spektraldarstellung einiger differentialoperatoren mit periodischen koeffizienten im raume der fastperiodischen funktionen *Stud. Math.* **25** 33–64
- [18] Chojnacki W 1991 Eigenvalues of almost periodic Schrödinger operators in $L^2(b\mathbb{R})$ are at most double *Lett. Math. Phys.* **22** 7–10
- [19] Krupa A and Zawisza B 1987 Ultrapowers of unbounded self-adjoint operators *Stud. Math.* **85** 107–23
- [20] Chalbaud E, Gallinar J-P and Mata G 1986 The quantum harmonic oscillator on a lattice *J. Phys. A: Math. Gen.* **19** L385–90
- [21] Abramowitz M and Stegun I A 1972 *Handbook of Mathematical Functions* (New York: Dover)
- [22] Avron J and Simon B 1981 The asymptotics of the gap in the Mathieu equation *Ann. Phys.* **134** 76–84
- [23] Finkel F, González-López A and Rodríguez M A 2000 A new algebraization of the Lamé equation *J. Phys. A: Math. Gen.* **33** 1519–42 (arXiv:math-ph/9908002)
- [24] Wightman A S 1995 Superselection rules: old and new *Il Nuovo Cimento B* **110** 751–69
- [25] Laddha A and Varadarajan M 2008 Polymer parametrised field theory *Phys. Rev. D* **78** 044008 (arXiv:0805.0208 [gr-qc])
- [26] Laddha A and Varadarajan M 2010 Polymer quantization of the free scalar field and its classical limit arXiv:1001.3505v1 [gr-qc]
- [27] Ashtekar A, Baez J, Corichi A and Krasnov K 1998 Quantum geometry and black hole entropy *Phys. Rev. Lett.* **80** 904 (arXiv:gr-qc/9710007 [gr-qc])
- [28] Agullo I, Barbero G J F, Diaz-Polo J, Fernandez-Borja E and Villaseñor E J S 2008 Black hole state counting in loop quantum gravity: a number theoretical approach *Phys. Rev. Lett.* **100** 211301 (arXiv:0802.4077 [gr-qc])
- [29] Engle J, Perez A and Noui K 2010 Black hole entropy and $SU(2)$ Chern–Simons theory *Phys. Rev. Lett.* **105** 031302 (arXiv:0905.3168 [gr-qc])
- [30] Krasnov K V 1997 Counting surface states in the loop quantum gravity *Phys. Rev. D* **55** 3505 (arXiv:gr-qc/9603025)
- [31] Barbero G J F and Villaseñor E J S 2011 The thermodynamic limit and black hole entropy in the area ensemble *Class. Quantum Grav.* **28** 215014 (arXiv:1106.3179 [gr-qc])